

The Milky Way rotation curve in Horava - Lifshitz theory

V.F. Cardone^{1,2}, N. Radicella^{3,4}, M.L. Ruggiero^{3,4}, M. Capone^{4,5}

¹ *Dipartimento di Scienze e Tecnologie dell' Ambiente e del Territorio, Università degli Studi del Molise, Contrada Fonte Lappone, 86090 - Pesche (IS), Italy*

² *Dipartimento di Scienze Fisiche, Università degli Studi di Napoli "Federico II", Complesso Universitario di Monte Sant' Angelo, Edificio N, via Cinthia, 80126 - Napoli, Italy*

³ *Dipartimento di Fisica, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 - Torino, Italy*

⁴ *I.N.F.N. - Sezione di Torino, via Pietro Giuria 1, 10125 - Torino, Italy*

⁵ *Dipartimento di Matematica, Università degli Studi di Torino, Via Carlo Alberto 10, 10125 - Torino, Italy*

Accepted xxx, Received yyy, in original form zzz

ABSTRACT

The Horava - Lifshitz (HL) theory has recently attracted a lot of interest as a viable solution to some quantum gravity related problems and the presence of an effective cosmological constant able to drive the cosmic speed up. We show here that, in the weak field limit, the HL proposal leads to a modification of the gravitational potential because of two additive terms (scaling respectively as r^2 and r^{-4}) to the Newtonian $1/r$ potential. We then derive a general expression to compute the rotation curve of an extended system under the assumption that the mass density only depends on the cylindrical coordinates (R, z) showing that the HL modification induces a dependence of the circular velocity on the mass function which is a new feature of the theory. As a first exploratory analysis, we then try fitting the Milky Way rotation curve using its visible components only in order to see whether the HL modified potential can be an alternative to the dark matter framework. This turns out not to be the case so that we argue that dark matter is still needed, but the amount of dark matter and the dark halo density profile have to be revised according to the new HL potential.

Key words: gravitation – dark matter – the Galaxy: kinematics and dynamics – galaxies: kinematic and dynamics

1 INTRODUCTION

Inspired by the Lifshitz theory in condensed matter physics, Horava (2009a,b) has recently proposed a new theory of gravity based on an anisotropic scaling of space and time in the UV limit. Usually referred to as Horava - Lifshitz (hereafter, HL) theory, the HL proposal shows a reduced invariance, dubbed *foliation preserving diffeomorphism invariance*, which however reduces to the standard one in the IR limit where General Relativity is recovered. As an attractive feature, the HL theory turns out to be power counting renormalizable which has motivated the great interest in investigating with great detail its theoretical and cosmological aspects. It is worth remembering that, in its original formulation, two conditions were imposed in order to drive the choice of the field action. First, the *projectability condition* was supposed to hold true. Since time plays a fundamental role from the very beginning, it was assumed that the lapse function (defined when working in the ADM formulation of gravity) has to be a projectable function on the spacetime foliation, that is to say a function of time only. Second, in or-

der to reduce the number of independent terms entering the action, the principle of *detailed balance* was used. Unfortunately, it soon became clear that this second condition leads to problems in the low energy limit (Lu et al. 2009) thus motivating the search for modification of the original HL theory where the action breaks the detailed balance condition either softly (Kehagias & Sfetsos 2009; Lee et al. 2009; Capasso & Polychronakos 2009; Kiristis & Kofinas 2009a; Kiristis & Kofinas 2009b) or not. In particular, Sotiriou, Visser and Weinfurtner (2009a,b) have worked out a modified HL theory with no detailed balance condition imposing that only parity preserving operators enter the potential.

Although the discussion about the foundations and the possible conceptual and phenomenological problems of the HL theory and its modified versions is still open, it is nevertheless worth systematically investigating its consequences at every scale. In particular, it is interesting to study its static spherically symmetric solutions since one can thus derive the gravitational potential generated from a point mass source. Recently, this problem has been addressed by Tang

& Chen (2009) for the HL theory with the projectability condition and no detailed balance. They argue that only the Minkowski or de Sitter spacetime are solutions, but we will show here that this is actually not the case. As a consequence, we find that the gravitational potential generated by a point mass differs from the Newtonian one because of the presence of additional terms depending on the HL coupling parameters. Such terms have to be taken into account when computing the potential generated by an extended system, such as a galaxy. Therefore, we work out a general formalism to estimate the rotation curve (i.e., the circular velocity v_c as function of the distance R from the centre) for an extended source showing that the HL theory can boost the $v_c(R)$ with respect to the Newtonian value. Motivated by this consideration, we then try to fit the Milky Way rotation curve using visible matter only to both constrain the HL parameters and investigating whether it can work as an effective dark matter component.

The plan of the paper is as follows. In Sect. 2, we look for static spherically symmetric solutions for the HL theory with projectability condition and show that there is indeed a new solution leading to a modified gravitational potential. A general formalism to compute the rotation curve for an extended system is presented in Sect. 3 and then used in Sect. 4 to work out the predicted rotation curve for the Milky Way. Here, we also present the data and the results of fitting them with our modified potential and no dark matter. Conclusions are finally given in Sect. 5.

2 THE POINT MASS POTENTIAL

The weak field limit of the HL theory as proposed by Sotiriou, Visser and Weinfurter has been yet discussed by Tang & Chen (2009, hereafter TC09) so that here we will only summarize the main steps and stress where our work differs from their one. As stressed in TC09, we start from the observation that not all the static spherically symmetric solutions found for (modifications of) HL theory preserve the projectability condition. In fact, if one assumes a metric

$$ds^2 = -N^2(r)dt^2 + \frac{dr^2}{g(r)} + r^2 d\Omega^2 \quad (1)$$

with $d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$ and use the coordinate transformation $dt = dt_{PG} - [(1 - N^2)^{1/2}/N^2]dr$, one can show that a necessary condition for the projectability condition to hold is that $g = N^2$. The line element therefore reads:

$$ds^2 = -N^2(t)dt^2 + \frac{(dr + N^r dt)(dr + N^r dt)}{f} + r^2 d\Omega^2. \quad (2)$$

The equations of motion for such a metric are given in TC09 and will not be repeated here for sake of shortness. The authors then argue that in the IR limit the f function is constrained to be 1 thus ending up with the usual Schwarzschild-de Sitter metric as only solution. We here show that this is actually not the case. To this aim, we first note that the Euler-Lagrange equation for $N_r = N^r/f$ in the IR limit reduces to

$$\frac{f'}{f} \frac{N_r}{r} = 0 \quad (3)$$

which is solved by either $N_r = 0$ or $f = \text{const.}$ The first choice gives Minkowski back when choosing $g_0 = 0$ and $f =$

1, or simply reduces to Minkowski by means of a redefinition of the radial coordinate. The second choice seems therefore more interesting. In such a case, the equations obtained by varying with respect to f and N reduce to

$$\frac{dN_r^2}{dr} + \frac{N_r^2}{r} + \frac{N^2(t)}{2f^2} \nu_1(r) = 0, \quad (4)$$

$$\int_0^\infty \left[\frac{dN_r^2}{dr} + \frac{N_r^2}{r} + \frac{N^2(t)}{2f^2} \nu_2(r) \right] r^3 dr = 0, \quad (5)$$

with

$$\begin{aligned} \nu_1(r) = & -g_0 \zeta^6 r + \frac{2\zeta^4(1-f)}{r} \\ & + \frac{2\zeta^2(1-f)}{r^3} [2g_2(1+7f) + g_3(1+5f)] \\ & + \frac{2(1-f)^2}{r^5} [4g_4(1+23f) + 2g_5(1+17f) \\ & + g_6(1+14f)] + \frac{8f(1-f)}{r^5} [2g_7(1+7f) \\ & + g_8(1-4f)], \end{aligned} \quad (6)$$

$$\begin{aligned} \nu_2(r) = & -g_0 \zeta^6 r + \frac{2\zeta^4(1-f)}{r} \\ & + \frac{2\zeta^2(1-f)^2}{r^3} (2g_2 + 4g_3) \\ & + \frac{2(1-f)^3}{r^5} (4g_4 + 2g_5 + g_6) \\ & + \frac{8f(1-f)^2}{r^5} (g_7 + g_8). \end{aligned} \quad (7)$$

For $f = 1$, we get $\nu_1(r) = \nu_2(r) = -g_0 \zeta^6 r$ so that Eqs.(4) and (5) are equal hence the solution for $N_r(r)$ is the same. However, differently from what stated in TC09, this is not the only possibility. Indeed, in order to have the same $N_r(r)$ solving both Eqs.(4) and (5), one must have $\nu_1(r) = \nu_2(r)$ which is possible by equating the coefficients of the terms with equal orders in r in the two functions. Comparing $\nu_1(r)$ and $\nu_2(r)$, one has just to equate the coefficients of the terms in r^{-3} and r^{-5} thus obtaining the following two equations:

$$(1-f) [2g_2(1+7f) + g_3(1+5f)] = (1-f)^2 (2g_2 + 4g_3), \quad (8)$$

$$\begin{aligned} 2(1-f)^2 [4g_4(1+23f) + 2g_5(1+17f) + g_6(1+14f)] + \\ 8f(1-f) [2g_7(1+7f) + g_8(1-4f)] = \\ 2(1-f)^3 (4g_4 + 2g_5 + g_6) + 8f(1-f)^2 (g_7 + g_8), \end{aligned} \quad (9)$$

which allows us to set two of the quantities ($g_1, \dots, g_8, f$) as a function of the others. Provided this condition has been satisfied, the solution of Eq.(4) automatically solves also the integral condition (5) so that we can limit our attention only to Eq.(4). This is a linear first order equation in $N_r^2(r)$ which can be analytically solved giving:

$$N_r(r) = \pm \sqrt{\frac{\mathcal{M}}{r^2} - \frac{N^2}{2f^2} \left(\frac{A}{3} r^2 + B - \frac{C}{r^2} - \frac{D}{3r^4} \right)} \quad (10)$$

where $\mathcal{M}$ is an integration constant and we have defined*

* Note that, following TC09, we have used units in which $Z = 1$ with Z a dimensional parameter of the HL theory. In order to

$$\begin{cases} A &= -g_0\zeta^6 \\ B &= 2\zeta^4(1-f) \\ C &= 2\zeta^2(1-f)[2g_2(1+7f) + g_3(1+5f)] \\ D &= 2(1-f)^2[4g_4(1+23f) + 2g_5(1+17f) \\ &\quad + g_6(1+14f)] + 8(1-f)f[2g_7(1+7f) \\ &\quad + g_8(1-4f)] \end{cases} \quad (11)$$

In order to derive the gravitational potential, we have first to rewrite the line element in the usual Schwarzschild-like form, i.e.

$$ds^2 = -g_{00}(r)dt^2 + g_{rr}(r)dr^2 + r^2d\Omega^2$$

and then use the general relations:

$$g_{00}(r) = N^2 - fN_r^2, \quad g_{rr}(r) = \frac{N^2}{f(N^2 - fN_r^2)}.$$

Fixing $N = 1$ (which is always possible by a rescaling of the t coordinate), we then get:

$$g_{00}(r) = 1 - \frac{f\mathcal{M}}{r} + \frac{1}{2f} \left(\frac{A}{3}r^2 + B - \frac{C}{r^2} - \frac{D}{3r^4} \right), \quad (12)$$

$$g_{rr}(r) = 1/fg_{00}(r). \quad (13)$$

Since the gravitational potential generated by a pointlike mass particle may be easily recovered from the usual relation $g_{00}(r) = 1 + 2\Phi(r)$, we then find:

$$\Phi(r) = -\frac{f\mathcal{M}}{2r} + \frac{1}{4f} \left(\frac{A}{3}r^2 + B - \frac{C}{r^2} - \frac{D}{3r^4} \right). \quad (14)$$

Note that, up to now, the two integration constants f and $\mathcal{M}$ are still undetermined. However, in order to recover the usual Newtonian potential in the GR limit (i.e., for $A = B = C = D = 0$), we must set $f\mathcal{M} = 2Gm$ with m the mass of the gravitational field source. It is worth noting that, for $m = 0$, the potential (14) leads to unphysical divergences in the metric because of the terms in C/r^{-2} and $D/3r^4$. In order to avoid this problem, we must impose that both C and D vanish for $m = 0$. The first trivial choice is to impose $C = D = 0$ identically so that one recovers the usual Schwarzschild-de Sitter case. As a more attractive possibility, one can postulate a dependence of the metric coefficient f on the mass. In such a case, Eqs.(8) and (9) can not be read as a relation among f and the HL couplings g_i since this will induce a dependence of the HL Lagrangian on the mass which is not possible. As such, we consider them as algebraic equations for f so that, in order to be fulfilled, we must equate the terms with equal powers of f on the two sides. We then obtain:

$$\begin{cases} g_2 = g_3 = 0 \\ g_4 = -(36g_5 + 15g_6 + 4g_7)/96 \\ g_8 = (16/3)g_7 \end{cases}$$

have (A, B, C, D) expressed in the more common $c = 1$ units, one has simply to multiply each term in Eq.(11) by ζ^{-4} with ζ having, in these units, the dimension of a length.

which implies $C = 0$ and

$$D = \frac{(1-f)^2}{12} [12(1-f)g_5 - 9(1-f)g_6 + 4(1-153f)g_7].$$

With these conditions, we can finally write the gravitational potential generated by a points mass m as:

$$\Phi(r) = \Phi_N(r) + \Phi_{HL}(r) \quad (15)$$

where $\Phi_N(r) = -Gm/r$ is the Newtonian potential and

$$\Phi_{HL}(r) = \frac{A}{12f}r^2 + \frac{B}{4f} - \frac{D}{12f} \frac{1}{r^4} \quad (16)$$

is the correction due to HL theory. Is is worth noting that, while the second term is simply a constant having no impact on the dynamics, the other two terms have a simple relation with previous results in literature. Indeed, the first one plays the role of a cosmological constant thus remembering the Schwarzschild-de Sitter solution already found in TC09. On the other hand, the third term has the same asymptotical behaviour of the corrections to the Newtonian potential in the Kehagias-Sfetsos (2009, hereafter KS) static spherically-symmetric solution of HL gravity. Iorio & Ruggiero (2009) have shown that such corrections are proportional to m^2r^{-4} thus suggesting that D should actually be a function of the mass m generating the gravitational field, a point which we will come back to later.

The corrective term may be conveniently written as:

$$\Phi_{HL}(r) = -\frac{GM_\odot}{r_s} \left[-\left(\frac{\eta}{\eta_A} \right)^2 - \frac{Br_s}{4fGM_\odot} + \left(\frac{\eta}{\eta_D} \right)^{-4} \right] \quad (17)$$

with $M_\odot$ the Sun mass and r_s an arbitrary chosen reference radius introduced to define the dimensionless quantity $\eta = r/r_s$. In Eq.(17), we have finally defined the scaling radii:

$$\begin{cases} \eta_A &= \left(\frac{12fGM_\odot}{Ar_s} \right)^{1/2} \\ \eta_D &= \left(\frac{Dr_s^{-3}}{12fGM_\odot} \right)^{1/4} \end{cases} \quad (18)$$

which can be used as model parameters instead of the (A, D) coefficients. Some caveats are in order here, First, note that, in order to preserve the interpretation of $(r_A, r_D) = r_s \times (\eta_A, \eta_D)$ as physical radii, one has to postulate that the (A, D) parameters are positive quantities thus narrowing the space of the parameters (g_0, g_5, g_6, g_7, f) entering (A, D) . Should A or D be negative, we could nonetheless define (r_A, r_D) as above and accordingly change the sign of the corresponding term in the potential. For definiteness, we will both A and D are positive so that the potential is given by Eq.(17) without any sign change. We then remember that, in our scheme, f is a dimensionless function of $\mu = m/M_\odot$, but its functional dependence can not be obtained in any way. The only constraint we have is $f(\mu = 0) = 1$ so that $D = 0$ and we recover the Schwarzschild-de Sitter solution in accordance with the fact that g_0 (and hence A) play the role of a cosmological constant term. In order to parametrize our ignorance we can redefine the above scaling radii as:

$$\begin{cases} \eta_A &= \left(\frac{12f_\odot GM_\odot}{Ar_s} \right)^{1/2} \left[\frac{f(\mu)}{f_\odot} \right]^{1/2} \\ \eta_D &= \left(\frac{D_\odot r_s^{-3}}{12f_\odot GM_\odot} \right)^{1/4} \left[\frac{D(\mu)}{D_\odot} \frac{f_\odot}{f(\mu)} \right]^{1/4} \end{cases} \quad (19)$$

where quantities labelled with a $\odot$ are evaluated for $\mu = 1$. It is worth wondering whether some hint on the functional expression of $f(\mu)$ can be retrieved. Up to now, we have postulated $f = f(\mu)$ in order to get a mathematically viable solution other than the Schwarzschild-de Sitter one. It is, however, worth noticing that a dependence of f on the mass may also be physically motivated. Although the HL theory is obtained by modifying General Relativity, it is still true that the properties of the spacetime are determined by the source of the gravitational field. As such, one can expect that the corrective HL term to the Newtonian potential is still related to the only property characterizing the field, i.e. the source mass m . Since the (A, B, D) coefficients in Eq.(??) are related to the HL couplings (and hence are universal quantities), the only way to introduce a dependence of the solution on the source properties is to postulate $f = f(\mu)$. Although qualitative, this discussion shows that our mathematical assumption is actually deeply related to a physical motivation.

Finally, we note that the second term in Eq.(17) simply adds a constant to the potential which has no effect in any situation of interest so that we will henceforth neglect this term. Note that this by no way means that B can be set to zero. Indeed, $B = 0$ means $f = 1$ so that also D vanish and we go back to the Schwarzschild-de Sitter solution, while we are here interested in the more general case. We therefore assume $B \neq 0$, but nevertheless neglect its contribution hereafter because drops off from the derivation of the quantities we are interested in.

3 THE ROTATION CURVE

The modified gravitational potential derived above deviates from the Newtonian one because of the additive terms in Eq.(17). Depending on the values of the scaling radii (r_A, r_D) , we can have different situations. As a general remark, we note that, for $\eta \ll 1$, the last term in Eq.(17) increases the potential with respect to the Newtonian one. On the contrary, for $\eta \gg 1$, the first term boosts the potential making it deviating from the Keplerian fall off. When considering the rotation curve, $v_c(r) = rd\Phi/dr$, we thus get a circular velocity which may significantly differ from the Newtonian one being larger than the classical value both in the inner and outer regions. It is therefore worth wondering whether such deviations may help in fitting spiral galaxies rotation curves without the need of other mass than the visible one. That is to say, we are here interested in investigating whether the HL modifications to the potential can also play the role of an effective dark halo. As a preliminary remark, we stress that a similar analysis could in principle be made also for the KS solution. However, in that case, the corrections fade away as r^{-4} and there is no cosmological constant term. It is therefore expected that the correction to the rotation curve in the outer regions are negligible so

that we argue that the KS solution can not play the role of an effective dark halo.

To this end, we have first to derive an expression for the rotation curve of an extended system generalizing the procedure adopted in the Newtonian gravity framework. In that case, the circular velocity in the equatorial plane is given by $v_c^2(R) = Rd\Phi/dR|_{z=0}$, with Φ the total gravitational potential. Thanks to the superposition principle and the linearity of the point mass potential on the mass m , this latter is computed by adding the contribution from infinitesimally small mass elements and then transforming the sum into an integral over the mass distribution. Such a simple procedure can not be applied in the HL case since the corrective term Φ_{HL} does depend on m in a way that we do not explicitly know so that a nonlinear dependence can not be excluded a priori. In order to overcome this difficulty, we have developed an alternative procedure which can be actually applied to any kind of potential provided some general conditions hold.

As a starting point, let us denote by $F_p(m, r)$ the gravitational force (per unit of test mass particle) generated by a point mass m . Whatever is the dependence of F_p on m , it is always true that the total force due to N particles of mass m is the sum of the single forces. As such, taking the continuum limit, we can estimate the (magnitude of the) total force as :

$$F(\mathbf{r}) = \int n(m, \mathbf{r}') F_p(m, |\mathbf{r} - \mathbf{r}'|) dm dV \quad (20)$$

where the integral is over the full mass range and volume and $n(m, \mathbf{r})$ is the star mass function (hereafter *MF*), i.e. the number of stars in the volume element dV with mass between m and $m + dm$. Because of its definition, we have :

$$\int_{m_{min}}^{m_{max}} n(m, \mathbf{r}) m dm = \rho(\mathbf{r}) \quad (21)$$

with $\rho(\mathbf{r})$ the mass density. As usual in literature, we will adopt the factorization hypothesis thus writing $n(m, \mathbf{r}) = \psi(m)\tilde{\rho}(\mathbf{r})$ with $\psi(m)$ the local[†] MF and $\tilde{\rho} = \rho(\mathbf{r})/\rho_0$ with $\rho_0 = \rho(\mathbf{R}_0)$. Defining $\mu = m/M_\odot$, we have the following normalization condition for the local MF :

$$\int_{\mu_{min}}^{\mu_{max}} \mu \psi(\mu) d\mu = \rho_0/M_\odot^2$$

so that henceforth we use as local MF the quantity $\mathcal{N}_1\psi(\mu)$ with

$$\mathcal{N}_1 = \frac{\rho_0}{M_\odot^2} \left[\int_{\mu_{min}}^{\mu_{max}} \mu \psi(\mu) d\mu \right]^{-1}. \quad (22)$$

Let us now assume that the point mass gravitational force may be factorized as :

$$F_p(\mu, r) = \frac{GM_\odot}{r_s^2} f_\mu(\mu) f_r(r)$$

with f_μ and $f_r(r)$ dimensionless functions depending on the particular form of the point mass gravitational potential Φ_p . Remembering that $\mathbf{F}_p = -\nabla\Phi_p$, it is only a matter of algebra to show that it is :

[†] Note that the term *local* refers to the Solar neighborhood when the galaxy is the Milky Way. For external galaxies, by *local*, we mean the MF in the neighborhood of a suitably chosen reference radius R_0 .

$$f_\mu(\mu) = \mu, \quad f_r(\eta) = 1/\eta^2,$$

for the Newtonian potential. For the HL corrective potential, we can split it as the sum of two terms, i.e.

$$\Phi_{HL}(r, r_A, r_D) = \Phi_A(r, r_A) + \Phi_D(r, r_D),$$

and then obtain:

$$f_r = \left(\frac{2}{\eta_{A\odot}} \right) \left(\frac{\eta}{\eta_{A\odot}} \right), \quad f_\mu = \left[\frac{f(\mu)}{f_\odot} \right]^{-1},$$

for Φ_A and

$$f_r = \left(\frac{4}{\eta_{D\odot}} \right) \left(\frac{\eta}{\eta_{D\odot}} \right)^{-5}, \quad f_\mu = \frac{D(\mu)}{D_\odot} \frac{f_\odot}{f(\mu)},$$

for the Φ_D term. We now use cylindrical coordinates (R, θ, z) and the corresponding dimensionless variables (η, θ, ζ) (with $\zeta = z/r_s$) and rely on the factorization hypotheses for both the MF and the point mass force to finally get:

$$\begin{aligned} F(\mathbf{r}) &= G \rho_0 r_s \\ &\times \frac{\int_{\mu_{min}}^{\mu_{max}} f_\mu(\mu) \psi(\mu) d\mu}{\int_{\mu_{min}}^{\mu_{max}} \mu \psi(\mu) d\mu} \\ &\times \int_0^\infty \eta' d\eta' \int_{-\infty}^\infty d\zeta' \int_0^\pi f_r(\Delta) \tilde{\rho}(\eta', \theta', \zeta') d\theta' \end{aligned} \quad (23)$$

with the shorthand notation

$$\Delta = [\eta^2 + \eta'^2 - 2\eta\eta' \cos(\theta - \theta') + (\zeta - \zeta')^2]^{1/2}. \quad (24)$$

The circular velocity in the equatorial plane along the major axis (which is the quantity typically measured for spiral galaxies) will be simply $v_c^2(R) = RF(R, \theta = z = 0)$. Since we will be interested in axisymmetric systems, we can set $\tilde{\rho} = \tilde{\rho}(\eta, \zeta)$. Moreover, our systems will be spiral galaxies, hence made out of a spheroidal bulge and a circular disk, so that a convenient choice for the scaling radius r_s will be the disk scalelength R_d . Under these assumptions, the rotation curve may then be evaluated as:

$$\begin{aligned} v_c^2(R) &= G \rho_0 R_d^2 \eta \\ &\times \frac{\int_{\mu_{min}}^{\mu_{max}} f_\mu(\mu) \psi(\mu) d\mu}{\int_{\mu_{min}}^{\mu_{max}} \mu \psi(\mu) d\mu} \\ &\times \int_0^\infty \eta' d\eta' \int_{-\infty}^\infty \tilde{\rho}(\eta', \zeta') d\zeta' \int_0^\pi f_r(\Delta_0) d\theta' \end{aligned} \quad (25)$$

with

$$\Delta_0 = \Delta(\theta = \zeta = 0) = [\eta^2 + \eta'^2 - 2\eta\eta' \cos \theta' + \zeta'^2]^{1/2}. \quad (26)$$

It is worth noting that, except in very particular cases (e.g., the Newtonian potential for a spherically symmetric mass distribution), the integrals over the coordinates in Eq.(25) have to be evaluated numerically. For computational reasons, it is useful to exchange the order of integration and resort to the logarithmic variables $\lambda = \log \eta$ and $\omega = \log \zeta$ so that we get the following equivalent expression for the rotation curve:

$$v_c^2(R) = G \rho_0 R_d^2 (\ln 10)^2 \text{dex}(\lambda)$$

$$\begin{aligned} &\times \frac{\int_{\mu_{min}}^{\mu_{max}} f_\mu(\mu) \psi(\mu) d\mu}{\int_{\mu_{min}}^{\mu_{max}} \mu \psi(\mu) d\mu} \\ &\times \int_{-1}^1 \frac{\tilde{\mathcal{R}}(\lambda, \xi)}{\sqrt{1 - \xi^2}} d\xi \end{aligned} \quad (27)$$

with $\text{dex}(x) = 10^x$ and

$$\tilde{R} = \int_{-\infty}^\infty \text{dex}(2\lambda') d\lambda' \int_{-\infty}^\infty f_r(\Delta_0) \tilde{\rho}(\lambda', \omega') \text{dex}(\omega') d\omega'. \quad (28)$$

Eqs.(25) and (27) are fully general and can be used to compute the rotation curve provided the expression for $f_\mu(\mu)$ and $f_r(\eta)$ are given. As a consistency check, it is easy to show that, for the Newtonian potential, the term depending on the MF is identically unity so that Eq.(25) reduces to a simple rewriting of the standard result. For the HL term, we get a dependence on the MF through the multiplicative term on the second row of Eq.(25). It is worth stressing that the MF here only plays the role of scaling up or down the rotation curve. This is a consequence of the HL corrective term (17) not depending on m . As such, indeed, one has simply to sum the contribution of all the stars notwithstanding their mass and this is indeed what the MF term gives in the HL case. The total rotation curve for the HL theory will finally be given as:

$$v_c^2(R) = v_N^2(R) + v_{HL}^2(R) = v_N^2(R) + v_A^2(R) + v_D^2(R)$$

where one can use the literature results for the computation of the Newtonian term $v_N^2(R)$ and our general rule with $f_\mu = 1$ and $f_r(\eta)$ given above to estimate the HL contribution $v_{HL}^2(R)$. Note that, in order to evaluate $v_c^2(R)$, we need to know not only the mass density (as in the Newtonian only case), but also the local MF because of the HL term.

4 THE MILKY WAY ROTATION CURVE

The HL theory has been originally conceived as an attempt to solve some quantum gravity related problems, but has soon attracted a lot of interest even as an alternative model for dark energy because of the presence of a cosmological constant like term. We have here shown that also the gravitational potential is modified with respect to the standard Newtonian one so that it is worth wondering if the additional terms may help in reconciling the data on the rotation curves with what is predicted from the visible matter only. To this end, one should fit the rotation curves of many spiral galaxies and find out that the fit is indeed satisfactorily good and the model parameters (r_A, r_D) are the same for all the galaxies being related to the HL Lagrangian couplings and hence universal quantities. Such a task is actually quite complicated because of the need for the knowledge of the local MF which is not at all constrained in external galaxies. Moreover, what one observes for these systems is the surface brightness profile due to the two visible components, namely the bulge and the disk. What we need to evaluate the rotation curve is, however, the mass density so that, even assuming simple functional forms for the bulge and the disk, we still do not have any knowledge of the bulge and disk mass-to-light ratios thus adding more parameters (and hence severe degeneracies) to be determined. As a first preliminary test, we therefore limit our attention to

the Milky Way (hereafter MW) only. Our position within it allows us to determine the local MF (e.g., by star counts or converting the observed luminosity function into a MF through an empirically determined M/L ratio) thus reducing the uncertainties of the problem. Moreover, we also have direct determinations of the galactic parameters of interest so that the only unknown quantities are the HL parameters (r_A, r_D) thus strongly reducing the possibility of degeneracies. In the following, we first describe the MW mass models and the data on the rotation curve and then present the fitting procedure and the results.

4.1 The mass models

It is common to describe a spiral galaxies as the sum of a visible component (made out of stars and gas) embedded in a dark halo (mainly populated by cold dark matter particles). Lacking any definitive laboratory evidence for DM, the only reason why one has to include it in galaxy modelling is to fit the rotation curves data. Since this evidence implicitly assumes that the Newtonian potential theory is the correct one, there is actually no compelling reason why one should add a priori a dark halo to the visible components in a modified gravity framework as the current HL theory. We therefore model the MW as made out of visible matter only distributed in a spheroidal bulge[‡] and a thick disc.

We follow Dehnen & Binney (1998, hereafter DB98) describing the bulge as a truncated power-law model, i.e. the scaled mass density reads :

$$\tilde{\rho}_b(R, z) = \left(\frac{\varrho}{R_b}\right)^{-\gamma} \left(1 + \frac{\varrho}{R_b}\right)^{\gamma-\beta} \exp\left(-\frac{\varrho^2}{R_t^2}\right) \quad (29)$$

with $\varrho^2 = R^2 + z^2/q^2$. Fitting the model to the infrared photometric *COBE/DIRBE* data yields values for the model parameters, namely :

$$\beta = \gamma = 1.8 \quad , \quad q = 0.6 \quad , \quad R_b = 1 \text{ kpc} \quad , \quad R_t = 1.9 \text{ kpc} \quad .$$

The reference density is not determined from the photometry, but may be related to the total bulge mass M_b as :

$$\rho_{0,b} = \frac{M_b}{2\pi \int_0^\infty R dR \int_{-\infty}^\infty \tilde{\rho}(R, z) dz} \quad (30)$$

Dwek et al. (1995) has found $M_b = (1.3 \pm 0.3) \times 10^{10} M_\odot$ for the bulge mass so that we set M_b to its central value neglecting the measurement uncertainty. We have then to set the local MF for the bulge. Zoccali et al. (2000) have determined it from the luminosity function of lower main sequence stars finding :

$$\psi_b(\mu) \propto \mu^{-\beta}$$

with $\beta = -1.33 \pm 0.07$ for stars in the mass range (0.15, 1.0) $M_\odot$. We extend it to the brown dwarfs region and neglect the measurement uncertainty thus setting

[‡] Actually, there are different evidences that the bulge has triaxial structure. We nevertheless use a less detailed spheroidal model since the bulge contribute to the dynamics over the range probed by the data is much smaller than the disc one, independently on the gravitational theory adopted. Such a simplification allows us to use Eq.(25) without introducing any significative bias.

$(\beta, \mu_{min}, \mu_{max}) = (-1.33, 0.03, 1.0)$ as our bulge MF parameters.

While important in the inner regions, the bulge only plays a minor role in determining the circular velocity over the regions probed by the data. The dynamics is here dominated by the disc component that we model as a double exponential. The mass density then reads :

$$\tilde{\rho}_d(R, z) = \exp\left(-\frac{R}{R_d}\right) \exp\left(-\frac{|z|}{z_d}\right) \quad (31)$$

where we follow DB98 fixing :

$$R_d = \kappa_d R_0 \quad , \quad z_d = 0.18 \text{ kpc} \quad ,$$

with $\kappa_d = 0.30 \pm 0.05$ a scaling parameter and R_0 the Sun distance from the MW centre. We then follow Cardone & Sereno (2005) adopting $\kappa_d = 0.30$ and $R_0 = 8.5$ kpc as fiducial parameters. The disk reference density is related to the disk total mass as :

$$\rho_{0,d} = \frac{M_d}{4\pi R_d^2 z_d} \quad (32)$$

while the disk mass is estimated as

$$M_d = 2\pi R_d^2 \Sigma_\odot \exp(R_0/R_d) \quad (33)$$

with $\Sigma_\odot = 48 \pm 8 M_\odot/\text{pc}^2$ (Kuijken & Gilmore 1989) the disc surface density at the Sun position. A caveat is in order here. The measured value of $\Sigma_\odot$ refers to the total mass, both stars and gas. In principle, we should separate the two components computing the total disc mass using $\Sigma_\star$ instead of $\Sigma_\odot$, with $\Sigma_\star = \Sigma_\odot - \Sigma_{ISM}$ and $\Sigma_{ISM} = 14.5 M_\odot/\text{pc}^2$ (Olling & Merrifield 2001) the gas surface density. We should then add a further disc like component for the gas with its density profile. As a reasonable approximation, we can, however, assume that the gas follows the same density profile as the stellar disc with the same scalelength radius so that the total disc mass is indeed given by Eq.(33).

As a final ingredient, we need the local disc MF. This is determined quite accurately thanks to our position in the disc plane. We refer the reader to Chabrier (2003) and references therein for a detailed discussion of this issue motivating our choice of adopting the Kroupa MF, i.e. a power-law MF with slope β changing with the mass range as :

$$\beta = \begin{cases} 0.3 & 0.01 \leq \mu \leq 0.08 \\ 1.3 & 0.08 \leq \mu \leq 0.50 \\ 2.3 & 0.50 \leq \mu \leq 1.0 \\ 4.5 & 1.0 \leq \mu \leq 120.0 \end{cases} .$$

The four different range contributes to the local mass density according to the following percentages :

$$(f_1, f_2, f_3, f_4) = (7.39, 48.21, 29.22, 15.18)\%$$

so that we can compute the multiplicative term entering the rotation curve for the HL term.

4.2 The data

Although not ideal targets for our test because of the poor knowledge of their parameter and MF, external galaxies are better suited for the measurement of precise and extended

rotation curves. On the contrary, our position in the MW disc equatorial plane makes it difficult to observationally determine this quantity. Indeed, one has to characterize the full line of sight velocity distribution in order to correct the observed velocity for asymmetric drift and projection effects. Moreover, the distance to the tracer should be known with great accuracy not to bias in a dangerous way the estimated circular velocity. Notwithstanding these difficulties, the rotation curve in the outer regions have been measured relying on Cepheids and HII regions. We follow DB98 to estimate the circular velocity from the Cepheids data of Pont et al. (1997) and the HII molecular clouds sample of Brand & Blitz (1993) using their same selection criteria. This dataset probes the radial range $8.2 \leq R$ (kpc) ≤ 18.94 and is affected by a large scatter due to both measurement errors and the inhomogeneity of the data. In order to smooth the data without introducing any bias or spurious correlations, we use the local regression method (Loader 1999). Originally proposed by Cleveland (1979) and further developed by Cleveland and Devlin (1988), the local regression technique combines much of the simplicity of linear least squares regression with the flexibility of nonlinear regression. The basic idea relies on fitting simple models to localized subsets of the data to build up a function that describes the deterministic part of the variation in the data, point by point. Actually, one is not required to specify a global function of any form to fit a model to the data so that there is no ambiguity in the choice of the interpolating function. Indeed, at each point, a low degree polynomial is fit to a subset of the data containing only those points which are nearest to the point whose response is being estimated. The polynomial is fit using weighted least squares with a weight function which quickly decreases with the distance from the point where the model has to be recovered. We hence use this method[§] to smooth the sample points and cut data with $R > 14$ kpc since the local regression method becomes unreliable for these points because of the sparseness of the sample in this region. We finally give it away points with $S/N \leq (S/N)_{min}$ where the threshold signal-to-noise ratio has been set to $(S/N)_{min} \simeq 8$ for reasons explained below. We will refer to this sample as the DB data in the following.

Another possible tracer of the velocity field in the outer regions is represented by the Blue Horizontal Branch (BHB) stars. Photometry and spectra from the Sloan Digital Sky Survey (SDSS) for a sample of 2466 BHB stars allow to determine the rotation curve up to ~ 60 kpc as described in Xue et al. (2008, hereafter X08). In order to extract $v_c(R)$ taking care of all the possible projection effects and systematic biases (due to, e.g., the survey mask and targetting algorithm), X08 have relied on mock matched observations based on two different cosmological simulations of MW-like galaxies. As such, there are two different datasets labelled as N and S depending on the simulation adopted. We here use only the N dataset since it is in better agreement with the DB sample over the range of overlap. Note that this sample (referred to hereafter as the SDSS dataset) is made out of only 10 points obtained by binning the 2466 BHB stars in almost equally spaced radial bins covering the range (7.5, 55) kpc. The S/N is quite high with a median value

$(S/N)_{med} \simeq 8$ so that we set $(S/N)_{min} = (S/N)_{med}$ in order to give the same statistical weight to both datasets. As a final remark, we note that, while the DB data probe only the outer disc being $R/R_d \simeq 3.2 - 5.4$, the SDSS sample extends mainly in the halo region most of the data having $R/R_d > 4$ and extending up to $R/R_d \simeq 20$. As such, the two sample nicely complement each other and allow us to check whether the outer rotation curve may be reproduced without any dark matter contribution and, at the same time, still preserving the agreement with the data in the disc region.

4.3 Fitting procedure

In order to constrain the HL model parameters, we employ a standard Bayesian approach first defining the likelihood function as :

$$\begin{aligned} \mathcal{L}(\mathbf{p}) &\propto \exp \left[-\frac{\chi^2(\mathbf{p})}{2} \right] \\ &= \exp \left\{ -\frac{1}{2} \sum \left[\frac{v_c^{obs}(R_i) - v_c^{th}(R_i)}{\varepsilon_i} \right]^2 \right\} \end{aligned} \quad (34)$$

where $\mathbf{p} = (\log \eta_A, \log \eta_C, \log \eta_D)$ are the HL model parameters[¶], $v_c^{obs}(R_i)$ and $v_c^{th}(R_i)$ are the observed (with a measurement error ε_i) and theoretically predicted values of the circular velocity at the radius R_i of the i -th point and the sum runs over the 101 DB and 10 SDSS data points.

The best fit is obtained by maximizing the likelihood $\mathcal{L}(\mathbf{p})$, but it is worth stressing that, according to the Bayesian philosophy, the best estimate of the parameter p_i is not the best fit one. On the contrary, one has to marginalize over the remaining parameters and look at the shape of the marginalized likelihood function defined as :

$$\mathcal{L}_i(p_i) \propto \int \mathcal{L}(\mathbf{p}) dp_1 \dots dp_{i-1} dp_{i+1} \dots dp_n$$

with n the total number of parameters. Actually, what we do is running a Monte Carlo Markov Chain code to efficiently explore the three dimensional parameter space $(\log \eta_A, \log \eta_C, \log \eta_D)$ and use of the histogram of the values for the parameter p_i to estimate the mean, the median and the 68 and 95% confidence ranges. Note that, because of degeneracies among the model parameters, the best fit parameters $\mathbf{p}_{bf}$ may also differ from the maximum likelihood ones $\mathbf{p}_{ML}$, i.e. the set obtained by maximizing each of the marginalized likelihood functions.

4.4 Results

As a preliminary discussion, it is worth clearly stating which are the parameters we can constrain. Eq.(25 shows that, in order to compute the HL contributions to the rotation

[§] See, e.g., Capozziello et al. 2007 for a step-by-step description.

[¶] We use logarithmic units in order to explore a wider range. Note also that, rigorously speaking, Eq.(34) is not correct since the SDSS data points are somewhat correlated being obtained by a binning procedure. However, since the bin spacing is quite large, it is likely that the correlation matrix (not available) is close to diagonal so that Eq.(34) is essentially correct.




Figure 1. Best fit HL model rotation curve superimposed to the DB and SDSS data.

curve, namely $v_A^2(R)$ and $v_D^2(R)$, we must know the functional expression of $f(\mu)$ entering $f_\mu(\mu)$ through the scaling radii (r_A, r_D) . Since we do not know this function, we can not separately constrain the parameters $(r_{A\odot}, r_{D\odot})$. It is, however, easy to show that:

$$\kappa_A v_A^2(R, \eta_{A,\odot}) = v_A^2(R, \eta_{A1}) ,$$

$$\kappa_D v_D^2(R, \eta_{D,\odot}) = v_D^2(R, \eta_{D1}) ,$$

where we have defined:

$$\kappa_A = \frac{\int_{\mu_{min}}^{\mu_{max}} [f(\mu)/f_\odot]^{-1} \psi(\mu) d\mu}{\int_{\mu_{min}}^{\mu_{max}} \psi(\mu) \mu d\mu} ,$$

$$\kappa_D = \frac{\int_{\mu_{min}}^{\mu_{max}} [D(\mu)/D_\odot] [f(\mu)/f_\odot]^{-1} \psi(\mu) d\mu}{\int_{\mu_{min}}^{\mu_{max}} \psi(\mu) \mu d\mu} ,$$

while $\eta_A = r_A/R_{eff}$ and $\eta_D = r_D/R_{eff}$ are considered as constant parameter not depending on μ and we assume that $v_A^2(R, \eta_A)$ and $v_D^2(R, \eta_D)$ are evaluated setting $f_\mu = \mu$ in Eq.(25). Our fit will then give constraints on (η_A, η_D) , while those on $(\eta_{A\odot}, \eta_{D\odot})$ could be derived provided a theoretically motivated functional expression for $f(\mu)$ is given.

Skipping to logarithmic units to investigate a larger range, we run a single chain with 150000 points reduced to ~ 10000 after cutting out the initial burn in phase and thinning the chain (with a step of 10) to avoid spurious correlations. The best fit point turns out to be:

$$(\log \eta_A, \log \eta_D) = (1.051, -0.060)$$

giving $\chi^2/d.o.f. = 523.56/109 = 4.80$. Needless to say, such a high reduced χ^2 is a strong evidence against the HL model with no dark matter. A simple look at Fig.1 shows that the large χ^2 is due mainly to the SDSS data. Indeed, it is $\chi_{DB}^2/d.o.f. = 235.91/99 = 2.38$ and $\chi_{SDSS}^2/d.o.f. = 287.65/8 = 35.96$ for the DB and SDSS samples, respectively. Although the model is clearly at odds with the data, we give below, for completeness, the constraints on the parameters from the analysis of their histograms along the chain. For $\log \eta_A$ we get:

$$\langle \log \eta_A \rangle = 1.049 \quad , \quad (\log \eta_A)_{med} = 1.050 \quad ,$$

$$68\% : (1.040, 1.059) \quad , \quad 95\% : (1.028, 1.066) \quad , \quad ,$$

while it is:

$$\langle \log \eta_D \rangle = -0.061 \quad , \quad (\log \eta_D)_{med} = -0.060 \quad ,$$

$$68\% : (-0.064, -0.056) \quad , \quad 95\% : (-0.068, -0.054) \quad ,$$

for $\log \eta_D$. Converting median values in linear units, we get:

$$(r_A, r_D) = (11.2, 0.87) \text{ kpc}$$

which allows us to make some qualitative interpretation of the results. Considering the form of the potential for the point mass case, we see that only the first term can actually boost the stellar contribution to the rotation curve. This term essentially translates in a contribution to $v_c(R)$ which linearly increases with R/r_A so that r_A must be small in order to fit the data in the DB probed region without dark matter. On the other hand, a too small value may lead to a $v_c(R)$ soon diverging thus spoiling down the agreement with the SDSS data. The value $r_A = 11.2$ kpc we find is the result of this compromise, but it is nevertheless unable to reconcile the model with the SDSS data. A similar compromise drives the fit in the estimates of $\log \eta_D$. Because of the scaling with r^{-4} in the point mass potential, the term entering r_D may contribute to the outer region circular velocity only if r_D is quite large. But, in such a case, R/r_D will be much smaller than 1 in the inner regions thus making the contribution of this term to overcome by orders of magnitude the Newtonian one in the regions where this latter is yet able to fit the data. As a consequence, r_D must be small so that it does not contribute enough to boost v_c with respect to the Newtonian value hence motivating the need for a not too large r_A . Indeed, we find an almost linear correlation between $\log \eta_A$ and $\log \eta_D$ along the chain motivated by the fact that the larger is r_D , the larger is the contribution to $v_c(R)$ of the r^{-4} term of the potential and hence the smaller is the need for the r^2 term which translates in a larger r_A .

5 CONCLUSIONS

Initially motivated by its attractive features from the point of view of quantum gravity, the HL proposal has soon become one of the most investigated theories of gravity. We have here complemented recent works on its cosmological consequences by addressing its impact on the gravitational potential. Contrary to the claim in TC09, we have demonstrated that static spherically symmetric solutions other than the Schwarzschild-de Sitter one exist. As a consequence, we have found a modified gravitational potential made out of the Newtonian $1/r$ one corrected by the addition of three further terms scaling as r^2 , which corresponds to the effect of a cosmological constant, and a quickly decreasing term, proportional to $1/r^4$. The importance of these two terms is parametrized in terms of two conveniently defined scaling radii, namely (r_A, r_D) , related to the couplings entering the HL Lagrangian. In order to consider astrophysically interesting situations, we have then developed a general formalism to compute the circular velocity curve provided the mass density and the mass function of the system are given. As an application, we have then evaluated the Milky

Way rotation curve using as only source of the gravitational field a spheroidal truncated power-law bulge and a double exponential disc. It turns out that the modified rotation curve is unable to fit the data thus demonstrating that the HL theory can not play the role of an alternative solution to the missing mass problem.

It is worth noting that Mukohyama (2009) has shown that classical solutions to the infrared (IR) limit of the HL theory can mimic General Relativity plus cold dark matter so that one can argue that it should be possible for the HL theory to provide an effective dark matter halo in apparent contradiction with our finding. Actually, this is not the case. First, Mukohyama only refers to the HL IR limit so that the Lagrangian he has considered does not include any potential term, while we here explicitly include this term through the coupling parameters g_i . Second, the cold dark matter term comes out as a consequence of the global Hamiltonian constraint being less restrictive than the local one typically imposed in General Relativity. In our static spherically symmetric metric, the global Hamiltonian constraint reduces to Eq.(5). In order to find our solution, we then impose that the integrand in Eq.(4) identically vanishes so that we are actually converting the global constraint in a local one thus going back to a situation similar to the General Relativity case. As a consequence, we argue that the possibility to get a matter term as an integration constant is lost in our approach.

The next step in our analysis of the HL theory on galactic scale should naturally be the inclusion of a dark halo. There are however some subtle issues making such a logical step forward not so easy to address. First, needless to say, there are few hints on which the dark halo mass density profile should be. All the models used in literature are motivated by the outcome of numerical simulations, such as, e.g., the popular NFW (Navarro et al. 1997) and Einasto (Einasto 1965; Cardone et al. 2005), or evidences from rotation curve fitting, such as the isothermal sphere (Binney & Tremaine 1987) and the Burkert (Burkert 1995; Burkert & Salucci 2000; Borriello & Salucci 2001) profile. All the previous works implicitly assume the validity of the Newtonian potential so that they are no more valid in a modified framework as the one we are using here. As a consequence, we have therefore to explore a wide class of density profile able to mimic most of the models in literature to finally select the most empirically motivated one. Moreover, as Eq.(25) shows, we also need to know the dark matter mass function which is completely unknown. Should the dark matter be composed of pointlike particles all having the same mass, we can adopt a Dirac δ leaving the mass of the particle as an unknown or setting it according to some particle physics model. As a final consequence, the fit to the rotation curve should determine the three HL scaling parameters (r_A, r_D) and $\mathcal{N}_{DM}$ halo parameters so that severe degeneracies among these $\mathcal{N}_{DM}$ quantities may take place. In order to reduce them, a better way should be to consider external galaxies rotation curves assuming that the MF is the same as the MW one. In such a way, we could take advantage of the data probing the full radial range and not only the outer disc as done here with the MW sample. Moreover, such a test can also probe the universality of the scaling radii (r_A, r_D) thus providing a further mandatory test of the HL model.

As a final remark, we want to stress that, should such

an analysis be successful, one has still to address a different issue. Let us suppose that we have indeed well fitted the rotation curves of a large sample of spiral galaxies thus determining a halo model and the values of (r_A, r_D). One could then use Eqs.(11) and (18) to infer constraints on the HL coupling parameters ($g_0, \dots, g_8$). Although we have thus five constraints (the two radii plus the two conditions imposed in the derivation of the potential) so being unable to determine all the eight quantities ($g_0, \dots, g_8$), one could nevertheless try to see whether the allowed region of the parameter space is consistent with what is inferred from cosmological analyses (Dutta & Saridakis 2009) or Solar System tests (see e.g. Iorio & Ruggiero 2009 and references therein). It is worth noting that such an attempt should give a consistent picture of the HL framework on all physical scales thus allowing to draw a deeper insight into its viability.

Acknowledgements. We warmly thank Mauro Sereno for helpful comments on a preliminary version of this manuscript. MC is supported by Regione Piemonte and Università degli Studi di Torino.

REFERENCES

- Brand, J., Blitz, L. 1993, *A&A*, 275, 67
- Binney, J., Tremaine, S. 1987, *Galactic dynamics*, Princeton University Press, Princeton (USA)
- Borriello, A., Salucci P. 2001, *MNRAS*, 323, 285
- Burkert, A. 1995, *ApJ*, 447, L25
- Burkert A., Salucci, P. 2000, *ApJ*, 537, L9
- Capasso, D., Polychronakos A.P. 2009, preprint arXiv:0911.1535
- Capozziello, S., Cardone, V.F., Troisi, A. 2007, *MNRAS*, 375, 1423
- Cardone, V.F., Sereno, M. 2005, *A&A*, 438, 545
- Cardone, V.F., Piedipalumbo, E., Tortora, C. 2005, *MNRAS*, 358, 1325
- Chabrier, G. 2003, *PASP*, 115, 763
- Cleveland, W.S. 1979, *J. Amer. Stat. Ass.*, 74, 829
- Cleveland, W.S., Devlin, S.J. 1988, *J. Amer. Stat. Ass.*, 83, 596
- Dehnen, W., Binney, J. 1998, *MNRAS*, 294, 429 (DB98)
- Dutta, S., Saridakis, E.N. 2009, preprint arXiv:0911.1435
- Dwek, E., Arendt, R.G., Hauser, M.G., Kelsall, T., Lisse, C.M., Moseley, S.H., Silverberg, R.F., Sodroski, T.J., Weiland, J.L. 1995, *ApJ*, 445, 716
- Einasto, J. 1965, *Trudy Inst. Astrofiz. Alma-Ata*, 5, 87
- Horava, P. 2009a, *Phys. Rev. D*, 79, 084008
- Horava, P. 2009b, *Phys. Rev. Lett.*, 102, 161301
- Iorio, L., Ruggiero, M.L., 2009, preprint arXiv:0909.2562
- Kehagias, A., Sfstos, K., 2009, *Phys.Lett. B*, 678, 123
- Kiristis, E., Kofinas, G. 2009a, *Nucl. Phys. B*, 821, 467
- Kiristis, E., Kofinas, G. 2009b, preprint arXiv:0910.5487
- Kuijken, K., Gilmore, G. 1989, *MNRAS*, 239, 605
- Lee, H.W., Kim, Y.W., Myung, Y.S. 2009, preprint arXiv:0907.3568
- Loader, C. 1999, *Local regression and likelihood*, Springer - Verlag, New York
- Lu, H., Jianwei, M., Pope, C.N. 2009, *Phys.Rev.Lett.*, 103, 091301
- Mukohyama, S. 2009, *Phys. Rev. D*, 80, 064005
- Navarro, J.F., Frenk, C.S., White, S.D.M. 1997, *ApJ*, 490, 493
- Pont, F., Queloz, D., Bratschi, P., Mayor, M. 1997, *A&A*, 318, 416
- Olling, R.P., Merrifield, M.R. 2001, *MNRAS*, 326, 164
- Sotiriou, M., Visser, T.P., Weinfurtner, S. 2009, *Phys. Rev. Lett.*, 102, 251601
- Sotiriou, M., Visser, T.P., Weinfurtner, S. 2009, *JHEP*, 10, 033

- Tang, J.Z., Chen, B. 2009, preprint arXiv:0909.4127 (TC09)
- Xue, X.X., Rix, H.W., Zhao, G., Re Fiorentin, P., Naab, T. et al.
2008, ApJ, 684, 1143
- Zoccali, M., Cassisi, S., Frogel, J.A., Gould, A., Ortolani, S., Renzini, A., Rich, R.M., Stephens, A.W. 2000, ApJ, 530, 418